



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Commentary

Thromboprophylaxis for medical inpatients with coronavirus disease 2019

M. Dalager-Pedersen^{1,2,*}, J. Bodilsen^{1,2}¹ Department of Infectious Diseases, Aalborg University Hospital, Aalborg, Denmark² Department of Clinical Medicine, Aalborg University, Aalborg, Denmark

ARTICLE INFO

Article history:

Received 5 May 2020

Received in revised form

17 May 2020

Accepted 18 May 2020

Available online 23 May 2020

Editor: L. Leibovici

Severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) and the disease it causes, coronavirus disease 2019 (COVID-19), have put health-care and financial systems worldwide under tremendous pressure. At present, COVID-19 has afflicted millions globally and the death toll is rapidly rising. Awaiting the development of effective and safe vaccines and antiviral therapies, researchers are struggling to better understand the disease and optimize supportive treatment.

Individuals hospitalized with COVID-19 are often immobilized with serious respiratory failure and have a remarkable pro-coagulant biochemical profile with elevated D-dimer levels and hyperinflammation. This has raised concern about increased risk of venous thromboembolism (VTE). Only a few studies specifically address this complication in COVID-19 and risk estimates range from 1% in ward patients to 35% in intensive care units [1,2]. However, most of these studies are hampered by small sample sizes in selected study populations treated in intensive care units at tertiary-care facilities with very short and incomplete follow up. Moreover, some of the studies also include VTEs with uncertain significance for risk of death, such as asymptomatic VTEs diagnosed by screening and subsegmental pulmonary embolisms. In conjunction with high mortality in previous studies, cumulative incidence findings of VTE may be inflated. Still, recent VTE estimates have garnered considerable media attention and are

currently extrapolated to all hospitalized individuals with COVID-19, while data on potential major bleeding complications and risk–benefit of anticoagulant therapy in COVID-19 are sparse.

Two recent retrospective observational studies have evaluated the effect of anticoagulation (AC) on mortality in individuals hospitalized with COVID-19.

Paranjpe et al. studied 2773 individuals hospitalized within the Mount Sinai Health System, New York City [3]. Overall, 786 individuals received treatment-dose AC during their hospital course and median time from admission to AC initiation was 2 days (interquartile range 0–5 days). Study participants were followed from hospital admission (T0) until discharge, death, or end of study. The authors compared mortality among AC users versus non-users and found similar mortality (22.5% versus 22.8%). In a sub-analysis among individuals receiving mechanical ventilation, AC was associated with greater benefit (mortality 29.1% versus 67.2%). However, as T0 was date of admission and AC initiation was delayed, the authors introduced immortal person-time among AC users, thereby conferring an artificial survival advantage to the AC group. Immortal time bias (or survivor treatment selection bias) can occur in survival analyses where patients who live longer are more likely to receive treatment than patients who suffer an early death [4]. As an example, Kaplan–Meier survival curves in the paper by Paranjpe et al. give the false illusion of improved survival among AC users when in fact ~25% of AC users were not at risk of death until after day 5 and all non-users were at risk from day 0.

A frequently cited study by Tang et al. examined the effect of AC, primarily enoxaparin 40–60 mg daily, on 28-day mortality in 449 hospitalized individuals with severe COVID-19 [5]. Exclusion criteria included hospitalization for <7 days and the AC group was defined as receiving AC for ≥7 days. In the primary analysis, the authors found no effect of AC on mortality (30.3% versus 29.7%). However, among individuals with sepsis-induced coagulopathy and in those with more than six-fold elevated D-dimer, mortality was lower among the individuals treated with AC. The study by Tang et al. is also at risk of immortal time bias unless everyone in the AC group initiated therapy on the day of admission, which is unclear. Of greater relevance, neither VTE events nor bleeding risk was detailed and the generalizability was limited on account of the inclusion criteria, meaning that just 449 of 1786 screened patients

* Corresponding author. Department of Infectious Diseases, Aalborg University Hospital, Mølleparkvej 4, 9000 Aalborg, Denmark.

E-mail address: midp@rn.dk (M. Dalager-Pedersen).

were included in the analysis. The authors sensibly conclude that AC may not benefit unselected COVID-19 patients, but should be considered in certain high-risk patients, for example those with sepsis-induced coagulopathy and markedly elevated D-dimer.

Of note, recent evidence from Italy suggests that deep venous thrombosis is an infrequent occurrence during COVID-19 and that filling defects on computed tomography angiography may be related to local pulmonary thrombi, and not to embolism, in which case heparin therapy is of questionable benefit [6].

Heparin therapy and thromboprophylaxis with heparin for individuals with infection and medical inpatients in general remains controversial [7,8]. Previous studies on thromboprophylaxis with low-molecular-weight heparin (LMWH) have found limited effect on clinically relevant outcomes in hospitalized medical patients with a number-needed-to-treat of approximately 250 to prevent symptomatic pulmonary embolism and a similar number-needed-to-harm in the form of major bleeding, resulting in little or no net benefit [7]. Moreover, thromboprophylaxis with LMWH has never been shown to prevent death in hospitalized medical patients including those with severe infection [9]. Prolonged thromboprophylaxis has been considered of potential benefit, but AC extended beyond hospital discharge for medical illness was not found to have an effect on risk for symptomatic VTE or death [10].

Guidelines on thromboprophylaxis and AC therapy in COVID-19 are rapidly emerging with differing recommendations. A recent position paper endorsed by several international societies suggested VTE risk stratification for all individuals with COVID-19 and pharmacological VTE prophylaxis in many cases [11]. The International Society for Thrombosis and Haemostasis has pushed the case for thromboprophylaxis with LMWH to all patients hospitalized with COVID-19 [12]. Other authorities have suggested intermediate or therapeutic doses of LMWH for hospitalized patients and extended VTE prophylaxis for up to 45 days post-discharge [11].

Although the COVID-19 pandemic confers a strong incentive on the medical community to act, we must remain adherent to evidence-based medicine and ethical considerations before changing guidelines from common practice, especially in prophylactic treatment of individuals. Consequently, there is a need for high-quality observational studies to better detail the incidence of VTE and bleeding events in individuals with COVID-19. There is also a need for information on risk factors and development of validated VTE and bleeding risk prediction models to identify those individuals with COVID who might benefit most from thromboprophylaxis. Even more importantly, we need well-conducted clinical trials on thromboprophylaxis in COVID-19 that explore clinically meaningful outcomes including symptomatic VTE, major bleeding events and death. These studies are needed to ensure that we do not harm patients, and may inform physicians and policy-makers of

the most efficient use of already heavily strained health-care resources.

Transparency declaration

The authors declare that there are no conflicts of interest. No external funding was received for this article.

Author contributions

MDP is lead and corresponding author. Conceptualization and investigation were by MDP and JB. The original draft was written by MDP and both MDP and JB contributed to reviewing and editing.

References

- [1] Klok FA, Kruip MJHA, van der Meer NJM, Arbous MS, Gommers DAMPJ, Kant KM, et al. Incidence of thrombotic complications in critically ill ICU patients with COVID-19. *Thromb Res* 2020. <https://doi.org/10.1016/j.thromres.2020.04.013>.
- [2] Goyal P, Choi JJ, Pinheiro LC, Schenck EJ, Chen R, Jabri A, et al. Clinical characteristics of Covid-19 in New York City. *N Engl J Med* 2020. <https://doi.org/10.1056/NEJMc2010419>.
- [3] Paranjpe I, Fuster V, Lala A, Russak A, Glicksberg BS, Levin MA, et al. Association of treatment dose anticoagulation with in-hospital survival among hospitalized patients with COVID-19. *J Am Coll Cardiol* 2020. <https://doi.org/10.1016/j.jacc.2020.05.001>.
- [4] Hanley JA, Foster BJ. Avoiding blunders involving “immortal time. *Int J Epidemiol* 2014;43:949–61.
- [5] Tang N, Bai H, Chen X, Gong J, Li D, Sun Z. Anticoagulant treatment is associated with decreased mortality in severe coronavirus disease 2019 patients with coagulopathy. *J Thromb Haemost* 2020;18:1094–9.
- [6] Cattaneo M, Bertinato EM, Biorchi S, Brizio C, Malavolta D, Manzoni M, et al. Pulmonary embolism or pulmonary thrombosis in COVID-19? Is the recommendation to use high-dose heparin for thromboprophylaxis justified? *Thromb Haemost* 2020. <https://doi.org/10.1055/s-0040-1712097>.
- [7] Lederle FA, Zylla D, MacDonald R, Wilt TJ. Venous Thromboembolism prophylaxis in hospitalized medical patients and those with stroke: a background review for an American College of Physicians Clinical Practice Guideline. *Ann Intern Med* 2011;155:602.
- [8] Zusman O, Paul M, Farbman L, Daitch V, Akayzen Y, Witberg G, et al. Venous thromboembolism prophylaxis with anticoagulation in septic patients: a prospective cohort study. *Qjm* 2015;108:197–204.
- [9] Kakkar AK, Cimminiello C, Goldhaber SZ, Parakh R, Wang C, Bergmann JF. Low-molecular-weight heparin and mortality in acutely ill medical patients. *N Engl J Med* 2011;365:2463–72.
- [10] Zayed Y, Kheiri B, Barbarawi M, Banifadel M, Abdalla A, Chahine A, et al. Extended duration of thromboprophylaxis for medically ill patients: a systematic review and meta-analysis of randomised controlled trials. *Intern Med J* 2020;50:192–9.
- [11] Bikdeli B, Madhavan MV, Jimenez D, Chuich T, Dreyfus I, Driggin E, et al. COVID-19 and thrombotic or thromboembolic disease: implications for prevention, antithrombotic therapy, and follow-up. *J Am Coll Cardiol* 2020. <https://doi.org/10.1016/j.jacc.2020.04.031>.
- [12] Thachil J, Tang N, Gando S, Falanga A, Cattaneo M, Levi M, et al. ISTH interim guidance on recognition and management of coagulopathy in COVID-19. *J Thromb Haemost* 2020;18:1023–6.